NUTRICHÈ

NURTURING LIFE

Head Office:
House No. 21, 2nd Floor, Khuzama Building, 2909 Abu Sufyan Bin Harb, Al Nzha Distt, Riyadh-KSA.
+966 50 514 1862

Corporate Headquarters:
13 Newstead Place, Halifax HX1 4TQ West Yorkshire UK.
+44 7471 972251

المكتب الرئيسى:
المنزل رقم ٢١، الطابق الثانى، مبنى خزامه، ٢٩٠٩ ابو سفيان بن حرب، النزهة ديست، الرياض-السعودية.
+٩٦٦ ٥٧ ٢٢٢ ٦٢٤٩

مقر الشركه:
١٣ نيوستيد بليس، هاليفاكس HX١ ٤TQ غرب يوركشاير، المملكة المتحدة.+٤٤ ٧٤٧١ ٩٧٢٢٥١



نيوتريش إي-سيلينيوم

NUTRICHÈ E-SELENIUM®

E-Selenium Immune Booster



Description

Nutrichè E-Selenium is a high-potency antioxidant and immune-enhancing supplement formulated for poultry, combining the synergistic benefits of Vitamin E (150,000 IU), Selenium (15,000 mg), Vitamin C (10,000 mg), Zinc (15,000 mg), and L-Lysine (10,000 mg). This formulation is designed to protect immune and muscle cells from oxidative stress, boost antibody production, support tissue regeneration, and enhance disease resistance during critical periods such as vaccination, infection, heat stress, and toxin exposure. By stabilizing cellular membranes, improving immune enzyme activity, and promoting protein synthesis, Nutrichè E-Selenium significantly improves fertility, hatchability, weight gain, FCR, and overall performance in broilers, layers, and breeders.

Composition

Vitamin E ----- 150,000 IU
Vitamin C ---- 10,000 mg
Selenium ---- 15,000 mg
Zinc ---- 15,000 mg
L-Lysine ---- 10,000 mg

Available SKU's

1 Liter Bottle (12 bottles per carton)
5 Liter Can (4 cans per carton)

Indications

Supports and maintains optimal fertility and hatchability in breeder flocks by protecting sperm and ova from oxidative damage, and improving reproductive tissue function under intensive production pressure.
For the prevention and treatment of deficiencies related to Vitamin E, Vitamin C, and Selenium, which may manifest as immunosuppression, oxidative muscle damage, or poor performance.
Highly effective in the management of muscular dystrophy, encephalomalacia, and pancreatic degeneration, particularly in rapidly growing or vitamin-deficient birds.
Recommended in clinical conditions such as infertility, subcutaneous edema, and feather abnormalities (e.g., ruffled feathers) resulting from vitamin-antioxidant imbalance or metabolic stress.
Significantly improves egg production and shell quality in laying and breeder hens by supporting cell membrane stability, ovarian response, and mineral metabolism.
Promotes efficient weight gain and feed conversion by improving nutrient utilization, immune efficiency, and tissue regeneration, especially during recovery from disease or post-vaccination stress.

DOSAGE & ADMINISTRATIONS

Preventive : 1 ml/10 lit of drinking water, once a week
Curative : 1 ml/6 lit of drinking water, for 3 days

Shake well before use.
Medicated drinking water should be used within 24 hours.

Storage Conditions

Store in a cool, dry place below 30°C
Protect from direct sunlight and moisture
Keep tightly closed in original container
Keep out of reach of children and unauthorized personnel